

1 **THE BIOCHAR OPTION TO IMPROVE PLANT YIELDS: FIRST RESULTS FROM**
2 **SOME FIELD AND POT EXPERIMENTS IN ITALY**

3
4

5 **Authors:** S. Baronti^(1*), G. Alberti⁽²⁾, G. Delle Vedove⁽²⁾, F. Di Gennaro⁽¹⁾, G. Fellet⁽²⁾, L.
6 Genesisio⁽¹⁾, F. Miglietta⁽¹⁾, A. Peressotti⁽²⁾, F.P. Vaccari⁽¹⁾

7
8
9

10
11

12 ⁽¹⁾ IBIMET-CNR Istituto di Biometeorologia, Consiglio Nazionale delle Ricerche
13 Via Giovanni Caproni, 8 - 50145, Firenze, Italy.

14
15

16 ⁽²⁾ Dipartimento di Scienze Agrarie ed Ambientali, Università di Udine
17 Via delle Scienze, 208 - 33100, Udine, Italy.

18

19 (*) Corresponding Author:

20 Silvia Baronti IBIMET-CNR, Via Caproni, 8 50145 Firenze, Italy

21 Tel. +390553033711

22 Fax +39055308910

23 Email: s.baronti@ibimet.cnr.it

1 **Abstract**

2 The pyrolysis conversion of agricultural residues into biochar and its incorporation in agricultural
3 soils, avoids CO₂ emissions providing a safe long term soil carbon sequestration. Furthermore,
4 biochar application to soil seems to increase nutrient stocks in the rooting zone, to reduce nutrient
5 leaching and to improve crop yields. This study highlights some preliminary results obtained using
6 biochar in two typical Italian agricultural crops. Two field experiments were made on durum wheat
7 (*Triticum durum* L.) in Central Italy and maize (*Zea mays* L.) in Northern Italy. In both the field
8 experiments, an increase in yields (+ 10% and + 6% in terms of grain production, respectively) was
9 detected after a biochar application of 10 t ha⁻¹. A further increase in grain production (+24%) was
10 detected when biochar is added with maize residues. The biochar dose effect curve was studied on
11 perennial ryegrass (*Lolium perenne* L.) in a pot experiment. The highest increase of dry matter
12 (+120 %) was obtained at a biochar rate of 60 t ha⁻¹ and above this threshold, a general reduction of
13 biomass was observed.

14

15 **Keywords**

16 Black carbon, crop yield, durum wheat, maize, sustainable agriculture

1 **Introduction**

2 Biochar is fine-grained and porous substance, similar in its appearance to charcoal produced by
3 natural burning. It is produced by pyrolysis, a thermo-chemical process where biomass is heated in
4 the absence of oxygen. As results, bio-oil, synthesis gas with different energy values and black
5 carbon (biochar) are obtained. Biochar can be used as soil amendment to improve soil quality, crop
6 yield and as a carbon (C) sequestration method. It may improve the physical structure of the soil
7 (Chan et al., 2007) and can also modify soil hydraulic properties (Chan et al., 2007; Gaskin et al.,
8 2007). Given that the pore size of biochar is relatively fixed, it increases available moisture in sandy
9 soils while has a neutral effect in medium textured soils and decreases moisture availability in clay
10 soils. In general, soil organic matter increases soil water holding capacity. In biochar-enriched *terra*
11 *preta* with their associated high levels of organic matter, Glaser et al. (2002) found a water retention
12 capacity that was 18% higher than in the adjacent soils. Biochar has been found to decrease nutrient
13 leaching thus enhancing nutrient availability (Chan et al., 2007; Yamato et al., 2006). Evidence
14 suggests that biochar porosity contributes to nutrient adsorption or covalent interaction on a large
15 surface area. Furthermore, its cation exchange capacity (CEC) is consistently higher than that of the
16 whole soil (Liang et al., 2006, Lehmann et al., 2003, Skjemstad et al, 1996; Lehmann et al., 2005).
17 In fact, the concentration of negative charges on biochar surfaces increases with age (Cheng et al.,
18 2006) as well as the adsorption of charged organic matter (Liang et al., 2006). According to
19 Lehmann et al. (2003), leaching of ammonium after biochar addition to soil was reduced by more
20 than 60% over four days of cropping rice compared to control treatments. Furthermore, these
21 authors measured a decrease in Ca, Mg and nitrate-N leaching.

22 Field experiments involving biochar application have been made for some crops and at different
23 locations, and positive yield responses have been often reported (Lehmann et al., 2003; Yamato et
24 al., 2006; Chan et al., 2007; Rondon et al., 2006; Rondon et al., 2007; Van Zwieten et al., 2008).
25 Most of these studies attributed the positive plant response to the effects of biochar on nutrients
26 availability (i.e. nutrient savings in terms of fertilizers or improved fertilizer use-efficiency). Some
27 studies (Yamato et al., 2006; Rondon et al., 2006; Van Zwieten et al., 2008) attributed the positive
28 plant response to the ability of biochar to increase or maintain the pH of soil, through liming.
29 However, most of studies made so far, have been conducted in tropical, semi-tropical and savannah
30 environments (Kimetu et al., 2008; Van Zwieten et al., 2008; Sinclair et al., 2008) while there is a
31 substantial lack of studies at mid-latitudes and in temperate climates.

32 Agricultural management may play a role in reducing the net greenhouse gas (UNFCCC, 2008) and
33 different agronomic strategies have been suggested and tested to reduce soil respiration rates and to
34 increase soil organic carbon. Those measures include the conversion of arable land to grasslands,

1 forest plantations and the implementation of crop management practices, such as conservation
2 tillage and the introduction of rotations based on nitrogen fixing crops (Lal and Bruce, 1999; Lal et
3 al., 1999; Lal, 2002). Recently, Lehmann et al. (2002) and Steiner et al. (2004) introduced the
4 concept of converting residues to biochar as an alternative agricultural method to reduce carbon
5 dioxide (CO₂) emissions. In fact, biochar can remain into soil for a long time (Glaser et al., 2001)
6 due to its stable structure and complex aromatic polycyclic form (Baldock and Smernik, 2002) thus
7 enhancing the resistance of C to microbial decay (Shindo, 1991; Cheng et al, 2008). Although
8 information about turnover time of biochar is scarce and mostly comes from short-term
9 decomposition experiments, it has been suggested that biochar stores atmospheric C from millennial
10 (Kuzyakov et al., 2009) to centennial timescales (Hamer et al., 2004). Based on those results,
11 biochar is currently considered as an interesting option to meet mitigation targets discussed by
12 United National Framework Convention on Climate Change (UNFCCC).

13 The objectives of the present study were to assess: 1) the effect of biochar doses on ryegrass
14 (*Lolium perenne* L.) production; 2) the effect of biochar application on durum wheat (*Triticum*
15 *durum* L.) and maize (*Zea mays* L.) yields; 3) the effect on maize yield of biochar application with
16 or without maize's crop residues.

17

18 **Materials and methods**

19 *Biochar*

20 Both field and pot experiments described in this paper were made using commercial horticultural
21 charcoal by Lakeland Coppice Products (England) obtained from coppiced woodlands (beech,
22 hazel, oak, birch). Biochar was prepared at pyrolysis temperatures of 500 °C in a transportable ring
23 kiln (215 cm in diameter and holding around 2 tons of hardwood). The material used was crushed
24 into particles smaller than 1 cm before application to the soil in order to increase the area: volume
25 ratio and to enhance its expected effects on soil properties.

26 C and N biochar contents were determined using a CHN Elemental Analyzer (Carlo Erba
27 Instruments, mod 1500 series 2). Samples of biochar were screened by means of a 2 mm sieve and
28 finally oven dried at 105 °C for 24 h. The dry samples were acid digested with a microwave oven
29 (CEM, MARSXpress) according to the EPA method 3052 (USEPA, 1995). The solutions obtained
30 after the mineralization were filtered (0.45µm PTFE) and diluted. Total contents of P, K, S, Ca, Mg
31 were determined by an ICP optical spectrometer (Varian Inc., Vista MPX) using scandium as
32 internal standard. The pH was measured in a soil/water solution at a 1/2.5 ratio. The main chemical
33 characteristics of the biochar used in the present study is reported on Table 1.

34

1 *Pot experiment*

2 In order to assess the dose-response curve of biochar, a pot experiment was made with perennial
3 ryegrass. Seeds were sown in pots of 5 l (0.2 m of diameter; 0.2 m of height) filled with a sandy-
4 loam soil at rate of 60 seeds per pot corresponding to 24 kg seeds ha⁻¹. Six application ratios of 0%,
5 0.3%, 0.8%, 1.7%, 2.8%, 3.3% (kg of biochar per kg of soil) were tested in a fully randomized
6 experimental design with three replicates. The biochar doses were equivalent to application rates of
7 0, 10, 30, 60, 100 and 120 t biochar ha⁻¹ assuming a soil bulk density of 1.2 g cm⁻³ up to a depth of
8 30 cm. After mixing, the pots were filled in order to ensure the same soil bulk density. Main soil
9 characteristics are reported in Table 2.

10 Biochar was added before sowing and the pots were regularly irrigated to prevent water stress, but
11 not fertilized. Aboveground biomass production was determined for two growth cycles (one and
12 two month after plant emergence) by manual clipping. Dry matter was determined on oven-dried
13 samples.

14

15 *Field experiments*

16 Two field experiments were performed in 2008. The first experiment was made in agricultural farm
17 near Empoli (Toscana, Central Italy) with durum wheat (cv. SOLEX). The field was located at an
18 altitude of 50 m a.s.l. with a sub-humid typical Mediterranean climate. According to long term
19 weather data, total rainfall was 750 mm and the mean annual temperature was 14 °C (ARSIA
20 Toscana, Italy). Main soil characteristics of the site are reported in Table 2.

21 A fully randomised experiment was made in 1.5 m² plots with two treatments (control C- and
22 biochar C+) and four replicates. Biochar addition corresponded to an overall rate of 10 t ha⁻¹ that
23 was incorporated in December 2007, an half (5 t ha⁻¹) immediately before sowing operation and an
24 half after germination. Wheat was sown on 17th December with a seeds density of 450 seeds m⁻². A
25 NP fertilizer was distributed at this time (22 kg N ha⁻¹ and 50 kg ha⁻¹ of P₂O₅) and a second nitrogen
26 fertilization using urea (92 kg N ha⁻¹) was made in February 2008. Plots were manually harvested
27 on 30th June 2008 in a central subplot of 0.7 m² to minimize edge effects. Total aboveground dry
28 matter was oven-dried at 105°C for 48 hours. Then, wheat ears were separated using a laboratory
29 thresher (LD 350, Wintersteiger, Ried, Austria). Nitrogen concentration of seeds was calculated by
30 Kjeldahl analysis.

31 The second field experiment was performed in Beano (Friuli Venezia Giulia, Italy) on a soil
32 cropped with irrigated continuous maize crop rotation. The mean annual temperature was 13.5° C
33 with a total rainfall of 1216 mm (OSMER Friuli Venezia Giulia).

34 The treatments were control (C-) and biochar addition (10 t ha⁻¹) with (CR) or without (C+)
35 residues of preceding corn crop. The three treatments were arranged in a randomized block

1 experiment with three replicates. Each experimental plot had an area of 20 m². On April 4th the cv
2 Kermess (from KWS maturity FAO class. 600) was sown at 7.2 seeds m⁻².
3 Fertilization was made at sowing adding 54 kg N ha⁻¹ and 138 kg P₂O₅ ha⁻¹ in the form of di-
4 ammonium phosphate. Urea was then added at two dates for a total of 359 kg ha⁻¹ of N (83 and 276
5 kg N ha⁻¹ on 09/05/2008 and 03/06/2008, respectively). The plots were weekly irrigated and no
6 water stresses were detected. Aboveground dry matter and yield were measured at harvest in a area
7 of 9 m² avoiding border effects of each plot. Main soil characteristics of the site are reported in
8 Table 2.

9

10 *Statistical analyses*

11 Treatment effects were analysed by analysis of variance (ANOVA) using SAS 9.1 (SAS Institute
12 Inc., Cary, NC, USA). The ANOVA included treatment C- and C+ in the durum wheat experiment
13 and C-, C+ and CR in the maize experiment. Since the experiment was a block design, block effect
14 was also included. Significant treatment effects (P<0.05) were explored further via a posteriori
15 treatment comparison using the Least-Squares means (LSMEANS) test with Bonferroni adjustment
16 for multiple comparison.

17

18 **Results**

19 *Pot experiment*

20 The higher was the rate of biochar added to the soil, the higher is the dry matter of ryegrass up to a
21 threshold of 60 t biochar ha⁻¹ (Figure 1). A significant difference among treatments was detected
22 (p=0.012). The ryegrass biomass increased by +29% and +120% at first clipping and +40% and
23 140% at second clipping in 0.8% of biochar (30 t ha⁻¹) and 1.7% (60 t ha⁻¹) treatments, respectively.
24 Germination was accelerated in the pots where biochar was added as likely consequence of increased
25 soil temperature (data non shown). Negative effects of biochar were observed above 1.7% of
26 biochar (60 t ha⁻¹) into the soil (60 t ha⁻¹).

27

28 *Field experiments*

29 In the field experiment with durum wheat, the phenological development of the wheat was not
30 affected by biochar addition. In fact, no differences were found between the treatments and no
31 major biotic and abiotic stresses were observed during the experimental period (data not shown).
32 The aboveground biomass was instead enhanced by biochar. The observed stimulation was 23%
33 respect to control with a value of 10.4 t ha⁻¹ and 8.6 t ha⁻¹ in C+ and C-, respectively (Figure 2;
34 p=0.054). Similarly, grain production was stimulated by 10% with a value 3.1 t ha⁻¹ and 2.4 t ha⁻¹ in

1 C+ and C-, respectively, but the significance of difference was low ($p=0.082$). The nitrogen content
2 of the wheat grains was not appreciably different between treated and un-treated plots (1.8% for
3 control and 1.84% for biochar treatment).

4 In the field experiment on maize, plant density at the end of the growing season was not
5 significantly different among treatments (data not shown). Aboveground biomass was also not
6 significant ($p=0.068$): 17.9, 22.5 and 25.8 t ha⁻¹ in C-, C+ and CR, respectively. Grain production,
7 instead, was significantly different ($p<0.01$) (Figure 3). In fact, a relative increase of grain yield in
8 C+ and CR in comparison to C- (+6% and +24% , respectively) was measured: 9.7, 10.3 and 12.1 t
9 ha⁻¹ in C-, C+ and CR, respectively. However, Bonferroni test showed a significant difference in
10 grain yield between C+ and CR ($P<0.05$) and between C- and CR ($P<0.01$), while no difference was
11 detected between C+ and C- ($P>0.05$).

12

13 **Discussion**

14 Agricultural productivity is often reported to increase with biochar application to soil, but always
15 not consistently. It is not yet clear what are the soil climatic conditions that favour increased yields,
16 and what are the most responsive species (Lehmann and Rondon, 2006).

17 Our results in the laboratory experiment showed that the rate of biochar application of 1.7%,
18 corresponding to 60 t ha⁻¹, caused the maximum stimulation in terms of dry matter production in
19 perennial ryegrass. Aboveground production stimulation by biochar addition was already observed
20 for grasses (Rondon et al., 2006; Lehmann et al., 2006) and this was interpreted as an overall
21 amelioration of growth condition. In our study increased availability of nitrogen was supposed to be
22 responsible for increased dry matter production (Gathorne-Hardy et al., 2009) as improved water
23 relations associated to biochar addition cannot be considered as the pots were maintained of the
24 field capacity throughout the experiment by means of ample irrigation. Above the 1.7% (60 t ha⁻¹)
25 threshold, the dry matter production of perennial ryegrass started to decline, probably because some
26 chemical and physical properties of the soil were modified by the high rate of biochar applied. A
27 possible explanation was given by Kishimoto and Sugiura, (1985) and Mikan and Abrams (1995)
28 who observed a general decrease of the biomass at high rate of the biochar application due to a
29 micronutrient deficiency induced by the increasing soil pH. Direct toxicity effect may also provide
30 an explanation, although these were not specifically investigated. Even a small biochar rate of 10 t
31 ha⁻¹ increased aboveground biomass (23%) and grain yield (10%) of durum wheat compared to
32 control. Comparable results were obtained by Steiner et al. (2007). A strict correlation between total
33 biomass production and grain yield is well known to exist in wheat as the harvest index is hardly
34 effect by different management and yield levels (Spaeth and Sinclair, 1995; Bindi et al., 1998). The

1 observed yield stimulation was likely due to a combination of improved soil water conditions
2 (Lehmann and Steiner, 2009), reduced nutrient leaching (Yanai et al., 2007) improved soil structure
3 and aggregate formation in the soil (Lehmann and Steiner, 2009). Increased nutrient availability
4 following biochar addition was observed in other field studies (Rondon et al., 2007) although such
5 effect cannot be directly detected in our experiment were the lack of difference between the N-
6 content of the grains of treated and untreated plants, supports this conclusion.

7 A positive effect on both aboveground biomass and grain was also observed in the field experiment
8 on maize. An increase of 26% and 6% in C+ and 44% and 24% in CR for aboveground biomass and
9 grain, respectively, was observed. A similar response was also obtained by Oguntunde et al. (2004).
10 The higher increase observed in CR in comparison to C+, even though not significant, is in
11 agreement with other studies (Glaser et al. 2002) and can be interpreted as a result of the positive
12 effect of maize stover residues of previous crop on the micro-scale physical properties of soil
13 aggregates, promoting their formation and stability (Blanco-Canqui and Lal, 2008).

14 Beneficial effects of biochar in terms of increased crop yield and improved soil quality have been
15 reported in several other studies (Iswaran et al., 1980; Glaser et al., 2002). However, Glaser et al.
16 (2002) underlined the fact that crop yields can be increased only if biochar is applied together with
17 inorganic or organic fertilizer as in the present study. In fact, Steiner et al. (2008) measured both a
18 higher soil N retention and an enhanced N cycling in fertilized plots that received biochar. This
19 could be due to a reduction in N leaching (higher retention of NH_4^+ ; N immobilization in microbial
20 biomass) or to a reduction of denitrification (Yanai et al., 2007) and we suppose that is the main
21 mechanism that can explain the observed yields increase in both our field experiments. However,
22 there could be other important interactions to explain the biochar stimulation of crop yield mostly
23 related to its higher stability in comparison to other organic amendments as well as the native soil
24 organic matter (SOM) (Steiner et al., 2007). Biochar can capture high amounts of exchange cations
25 (Lehmann et al., 2003) because of its high porosity and surface/volume ratio and can improve plant
26 nutrients uptake and P, Ca, K availability (Chan et al., 2007; Yamato et al., 2006). Biochar should
27 no longer seen as an inert material that remains unaltered in the soil where it is deposited. In fact,
28 during its permanence in soils, biochar is slowly oxidized, carboxylic groups are produced, cation
29 exchange capacity and oxygen carbon ratio on the biochar surface increases (Brodowski et al.,
30 2005), improving the capacity of biochar to retain nutrients in the long term. However, chemical
31 and physical characteristics of biochar depends on the nature of the feedstocks used and on the
32 operating conditions of pyrolysis process (Gundale and De Luca, 2006). Therefore, yield responses
33 are currently difficult to predict, and global patterns need to be identified to move towards an
34 understanding of the crop production potential using biochar.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

Conclusions

The results shown in this paper highlight a number of conclusions that can have practical importance for future experiment and real-world application of biochar in agriculture.

- A) *positive effect on growth an yield are sustained even at very high rates of biochar addition to the soil.* In the pot experiment, negative effects were observed when more than 1.7% (60 t ha⁻¹) of biochar were added. The preliminarily type of investigation and the limited prediction capacity of the pot experiment do not allow excessive generalization, but outline the fact that negative effects are unlikely under-realistic application scenarios. The addition of 30-60 t ha⁻¹ of biochar, to a 30 cm deep soil (1.2 bulk density) would in fact correspond to 0.6 -1.2% rate, below the 1.7% rate over which negative effect were observed.
- B) *even relatively small rates of biochar addition can have positive effects on crop growth and yield.* Although the limited scope of our initial trial does not allow to properly understand the mechanism underlying the observed increase in yield, there are convincing indication that biochar acts as a soil amendant capable of enhancing resource availability for the crop. Higher soil N-retention can be responsible for increased productivity (Steiner et al., 2008) but also enhanced N-cycling (reduction of denitrification) can contribute (Yanai et al., 2007). Increased soil water capacity, mechanical characteristics promoting soil organic matter stabilization, (Steiner et al., 2007), the availability of high amounts of exchange cations associated to high porosity of biochar (Lehmann et al., 2003) are additional factors that requires further specific investigation in field experiments.
- C) *enhanced yield and soil amelioration due to biochar application is always accompanied by long-term C-sequestration.* When plant residues are transformer into char by pyrolysis they reach very stable condition that can only be partly decomposed by soil micro-organisms. There is evidence in literature that the largest part carbon contained in the biochar can remain undecomposed for hundred years or even millennia.
- D) *the positive effects of biochar application, its potential for C-sequestration are accompanied by the production of thermal-energy during the pyrolysis.* When such energy is efficiently used it can replace the use of fossil fuels and therefore further contribute to an overall reduction of net emission of greenhouse gases to the atmosphere.

Our study must be seen as a first attempt to frame the research of biochar within a coherent scientific approach in Italy. More studies are definitely required to expand phenomenological observation and to improve or mechanistic understanding of processes.

1

2 **Acknowledgement**

3 Diego Chiabà (University of Udine) is acknowledged for valuable technical assistance during the
4 field experiment. This study contributes to the by the Italian Biochar Association (ICHAR

5 <http://www.ichar.org>)

1 **References**

- 2 Baldock J.A., Smernik R.J., 2002. Chemical composition and bioavailability of thermally altered
3 *Pinus resinosa* (Red pine) wood. *Org. Geochem.*, 33: 1093-1109
- 4 Bindi M., Sinclair T.R., Harrison J. 1998. Analysis of seed growth by increase in harvest index.
5 *Crop Science*, 39: 486-493
- 6 Blanco-Canqui H., Lal R. 2009. Corn Stover removal impacts on micro-scale soil physical
7 properties. *Geoderma*, 145: 335-346
- 8 Brodowski S., Amelung W., Haumaier L., Abetz C., Zec W. 2005. Morphological and chemical
9 properties of black carbon in physical soil fractions as revealed by scanning electron
10 microscopy and energy-dispersive X-ray spectroscopy. *Geoderma*, 128: 116-129
- 11 Chan K. Y., Van Zwieten L., Meszaros I., Downie A., Joseph S. 2007. Agronomic values of
12 greenwaste biochar as a soil amendment. *Aust. J. Soil Res.*, 45: 629-634
- 13 Cheng C. H., Lehmann J., Engelhard M.H. 2008. Natural oxidation of black carbon in soils:
14 Changes in molecular form and surface charge along a climosequence. *Geochim. Cosmochim.*
15 *Ac.*, 72: 1598-1610
- 16 Cheng C.H., Lehmann J., Thies J.E., Burton S.D., Engelhard M.H. 2006. Natural oxidation of black
17 carbon in soils: changes in molecular form and surface charge along a climo-sequence.
18 *Geochim. Cosmochim. Ac.*, 72: 1598-1610
- 19 Gaskin J.W., Speir A., Morris L.M., Ogden L., Harris K., Lee D., Das K.C. 2007. Potential for
20 pyrolysis char to affect soil moisture and nutrient status of loamy sand soil. In: Georgia Water
21 Resources Conference. University of Georgia.
- 22 Gathorne-Hardy A., Knight J., Woods J. 2009. Biochar as a soil amendant positively interacts with
23 nitrogen fertilizer to improve barley yields in the UK. *IOP Conf.Series: Earth and*
24 *Environmental Sciences* 6 doi10.1088/1755-1307/6/7/372052
- 25 Glaser B., Haumeier L., Guggenberger G., Zech W. 2001. The 'Terra Preta' Phenomenon: a model
26 for sustainable agriculture in the humid tropics. *Naturwissenschaften*, 88: 37-41
- 27 Glaser B., Lehmann J., Zech W. 2002. Ameliorating physical and chemical properties of highly
28 weathered soils in the tropics with charcoal a review. *Biol. Fert. Soils.*, 35: 219-230
- 29 Gundale M.J., De Luca T.H. 2006. Temperature and source material influence ecological attributes
30 of ponderosa pine and Douglas-fir charcoal. *Forest Ecol. Manag.*, 231: 86-93
- 31 Hamer U., Marschner B., Brodowski S., Amelung W. 2004. Interactive priming of blackcarbon and
32 glucose mineralization. *Org. Geochem.*, 35: 823-830
- 33 Iswaran V., Jauhri K.S., Sen A. 1980. Effect of charcoal, coal and peat on the yield of moong,
34 soybean and pea. *Soil Biol. Biochem.*, 12:191-192

- 1 Kimetu J.M., Lehmann J., Ngoze S.O., Mugendi D.N., Kinyangi J.M., Riha S., Verchot L., Recha
2 J.W., Pell A.N. 2008. Reversibility of soil productivity decline with organic matter of
3 differing quality along a degradation gradient. *Ecosystems* (published online).
4 doi:10.1007/s10021-008- 9154-z
- 5 Kishimoto S., Sugiura G. 1985. Charcoal as a soil conditioner. In: Symposium on forest products
6 research, *Int. Achieve. Future*, 5: 12-23
- 7 Kuzyakov Y., Subbotina I., Chen H., Bogomolova I., Xu X. 2009. Black carbon decomposition and
8 incorporation into soil microbial biomass estimated by ¹⁴C labeling. *Soil Biol. Biochem.*, 41:
9 210–219
- 10 Lal R. 2002. Carbon sequestration in dry land ecosystems of West Asia and North Africa. *Land*
11 *Degrad.Dev.*, 13: 45-59
- 12 Lal R., Bruce J.P. 1999. The potential of world cropland soils to sequester C and mitigate the
13 greenhouse effect. *Environ. Sci. Policy*, 2: 177-185
- 14 Lal R., Kimble J.M., Follet R.F., Cole C.V. 1999. The potential of U.S. cropland to sequester
15 carbon and mitigate the greenhouse effect. Lewis Publisher, New York, NY., p. 128
- 16 Lehmann J., Steiner J. 2009. *Biochar for Environmental Management*. published by Earthscan
17 Publishers Ltd (ISBN 978-1-84407-658-1) March 2009
- 18 Lehmann J., Rondon M. 2006. Bio-char soil management on highly weathered soils in the humid
19 tropics, in N. Uphoff (ed) *Biological Approaches to Sustainable Soil Systems*, CRC Press,
20 Boca Raton, FL, USA, pp 517-530
- 21 Lehmann J., Liang B., Solomon D., Lerotic M., Luizão F., Kinyangi J., Schäfer T., Wirick S.
22 Jacobsen C. 2005. Near-edge X-ray absorption fine structure (NEXAFS) spectroscopy for
23 mapping nano-scale distribution of organic carbon forms in soil: Application to black carbon
24 particles. *Global Biogeochem. Cl.*, 19: GB1013 DOI 10.1029/2004GB002435
- 25 Lehmann J., Da Silva J.P. jr., Steiner C., Nehls T., Zech W., Glaser B. 2003. Nutrient availability
26 and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin:
27 fertilizer, manure and charcoal amendments. *Plant Soil.*, 249: 343-357
- 28 Lehmann J., Da Silva J.P. jr., Rondon M., Cravo M.S., Greenwood J., Nehls T., Steiner C., Glaser
29 B. 2002. Slash-and-char a feasible alternative for soil fertility management in the central
30 Amazon?, 17th World Congress of Soil Science, Bangkok, Thailand, Paper No. 449
- 31 Liang B., Lehmann J., Solomon D., Kinyangi J., Grossman J., O'Neill B., Skjemstad J.O., Thies J.,
32 Luizao F.J., Petersen J., Neves E.G. 2006. Black Carbon increases cation exchange capacity in
33 soils. *Soil Sci. Soc. Am. J.*, 70: 1719-1730

- 1 Mikán C.J., Abrams M.D. 1985. Altered forest composition and soil properties of historic charcoal
2 hearths in southeastern Pennsylvania. *Can. J. of Forest. Res.*, 25: 687-696
- 3 Oguntunde P.G., Fosu M., Ajayi A.E., Van de Giesen N. 2004. Effects of charcoal production on
4 maize yield, chemical properties and texture of soil. *Biol. Fertil. Soils.*, 39: 295-299
- 5 Rondon M.A., Lehmann J., Ramírez J., Hurtado M. 2007. Biological nitrogen fixation by common
6 beans (*Phaseolus vulgaris* L.) increases with bio-char additions. *Biol. Fertil. Soils.*, 43: 699-
7 708
- 8 Rondon M., Molina D., Ramirez J., Amezquita E., Major J., Lehmann J. 2006. Enhancing the
9 productivity of crops and grasses while reducing greenhouse gas emissions through bio-char
10 amendments to unfertile tropical soils. Poster presented at the World Congress of Soil
11 Science, Philadelphia, PA, July 9-15
- 12 Rondon M., Ramirez A. and Hurtado M. 2004. Charcoal additions to high fertility ditches enhance
13 yields and quality of cash crops in Andean hillsides of Columbia, CIAT Annual Report Cali,
14 Colombia
- 15 Shindo H. 1991. Elementary composition, humus composition, and decomposition in soil of charred
16 grassland plants. *Soil Sci. Plant Nutr.*, 37: 651-657
- 17 Sinclair K., Slavich P., Van Zwieten L., Downie A. 2008. Productivity and nutrient availability on a
18 Ferrosol: biochar, lime and fertilizer, in Proceedings of the Australian Society of Agronomy
19 conference, 21-25th September 2008, Adelaide Australia, (in press)
- 20 Skjemstad J.O., Clarke P., Taylor J.A., Oades J.M., McClure S.G. 1996. The chemistry and nature
21 of protected carbon in soil. *Aus. J. Soil Res.*, 34: 251-271
- 22 Spaeth S.C., Sinclair T.R. 1985. Linear increase in soybean harvest index during seed-filling.
23 *Agron. J.*, 77: 207-211
- 24 Steiner C., Glaser B., Teixeira W.G., Lehmann J., Blum W.E.H., Zech W. 2008. Nitrogen retention
25 and plant uptake on a highly weathered central Amazonian Ferralsol amended with compost
26 and charcoal. *J. Plant Nutr. Soil Sci.*, 171: 893-899
- 27 Steiner C., Teixeira W.G., Lehmann J., Nehls T., Luis J., De Macedo L.V., Blum W.E.H., Zech W.
28 2007. Long term effects of manure, charcoal and mineral fertilization on crop production and
29 fertility on a highly weathered central Amazonian upland soil. *Plant Soil.*, 291: 275-290
- 30 Steiner C., Teixeira W.G., Lehmann J., Zech W. 2004. Microbial response to charcoal amendments
31 of highly weathered soils and Amazonian Dark Earths in Central Amazonia preliminary
32 results, in B. Glaser and W. I. Woods (eds) *Amazonian Dark Earths: Explorations in Time
33 and Space*, Springer, Berlin, Germany, pp 195-212

- 1 UNFCCC. 2008. Analysis of possible means to reach emission reduction targets and of relevant
2 methodological issues. FCCC/TP/2008/2, available at: <http://unfccc.int>
- 3 Van Zwieten L., Kimber S., Sinclair K., Chan K.Y., Downie A. 2008. Biochar: Potential for climate
4 change mitigation, improved yield and soil health, in New South Wales Grassland
5 Conference, 2008
- 6 Yamato M., Okimori Y., Wibowo I.F., Anshori S., Ogawa M. 2006. Effects of the application of
7 charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical
8 properties in South Sumatra, Indonesia. *Soil Sci. Plant Nutr.*, 52: 489-495.
- 9 Yanai Y., Toyota K., Okazaki M. 2007. Effects of charcoal addition on N₂O emissions from soil
10 resulting from rewetting air-dried soil in short-term laboratory experiments. *Soil Sc. Plant*
11 *Nutr.*, 53: 181-188

1 **Tables**

2

3 Table 1 – Characteristics of the biochar used in this study.

BIOCHAR CHARACTERISTICS

<i>Chemical Element</i>	<i>Value</i>
Total C (g kg ⁻¹)	840
Total N (g kg ⁻¹)	12
Available N (g kg ⁻¹)	0.03
P (g kg ⁻¹)	0.5
K (g kg ⁻¹)	4.3
Ca (g kg ⁻¹)	2.6
S (g kg ⁻¹)	1.1
Mg (g kg ⁻¹)	2.8
C:N	70
pH (1:2.5 H ₂ O)	7.2

4

5 Table 2 – Texture and chemical Characteristics of the experimental soils.

	Pot experiment	Empoli	Beano
Gravel (% of dry bulk soil) $\varnothing > 2\text{mm}$	n.a.	n.a.	32
Sand (g kg ⁻¹) ^a $2\text{ mm} > \varnothing > 0.05\text{mm}$	550	350	270
Silt (g kg ⁻¹) $0.05\text{ mm} > \varnothing > 0.002\text{mm}$	300	360	580
Clay (g kg ⁻¹) $\varnothing < 0.002\text{ mm}$	150	290	150
Bulk density (Mg m ⁻³)	1.6	n.a.	1.2
OC (g kg ⁻¹) ^b	n.a.	18	19
N (g kg ⁻¹) ^b	n.a.	1.6	2
CEC (meq/100g) ^b	n.a.	16	n.a.
pH ^b	6.7	8.0	7.1

6 ^a) it refers to fine (<2 mm) texture fraction.

7 ^b) Method used: CaCl₂ 0.01M, 1:2.5 CaCl

8

9

10

1 **Figure captions**

2

3 Fig. 1 Aboveground biomass of *Lolium perenne* in the laboratory experiment in the first and second
4 growth cycle at different rates of biochar ($t\ ha^{-1}$). Each values is the average of 3 replicates. Vertical
5 bars indicate +/- standard error. Different letters indicate a significant difference respect to control
6 (rate of biochar corresponding to 0) ($P<0.05$).

7


8 Fig.2 Aboveground biomass (open bars) and grain production (solid bars) of durum wheat in the
9 field experiment in Central Italy. C+ is the biochar, C- is the control, respectively. Each value is the
10 average of 4 replicates. Vertical bars indicate +/- standard error. Different letters indicate a
11 significant difference ($P<0.05$).

12

13 Fig. 3 Aboveground biomass (open bars) and grain production (solid bars) of maize in the field
14 experiment in North Italy. C+ is the biochar, C- is the control and CR is the biochar + residues,
15 respectively. Each value is the average of 3 replicates. Vertical bars indicate +/- standard error.
16 Different letters indicate a significant difference ($P<0.05$).

1 **Figure 1**

2




3

4

5


1 **Figure 2**

2



3

1 **Figure 3**



2

3